

Hip Fracture Deaths in Utah Hospitals (IQI 19)

Adults 18 Years and Over: 2005-2007

Hospital	Patients (Denominator)	Actual Rate	Expected Rate	Statistical Rating
Utah Overall	843	0.6%	0.8%	**
Cottonwood Hospital	41	0.0%	1.5%	**
Dixie Regional	53	0.0%	0.4%	**
LDS Hospital	108	0.9%	1.0%	**
McKay-Dee Hospital	55	3.6%	1.4%	**
St. Mark's	63	0.0%	0.4%	**
University Health Care	139	0.7%	0.7%	**
Utah Valley Regional	88	0.0%	1.1%	**

All Utah hospitals in this table had about the same rate of deaths as expected compared to similar patients in the U.S.

[See additional hospitals that treated at least one patient but less than 30 patients.](#)

[View Data Limitations.](#)

Rating: based on the Exact 95% Confidence Interval test of statistical significance.

*** Lower % deaths than expected, ** Same % deaths as expected, * Higher % deaths than expected

Actual deaths: percentage of patients who received care for this condition or procedure and died in this hospital.

Expected deaths: percentage of patients who were expected to die at this hospital if it performed the same as other U.S. hospitals that treated similar patients (age, gender, how ill the patients were).