

Hip Fracture Deaths in Utah Hospitals (IQI 19)

Adults 18 Years and Over: 2006-2008

Hospital	Patients (Denominator)	Actual Rate	Expected Rate	Statistical Rating
Utah Overall	869	0.6%	0.7%	**
Dixie Regional	58	0.0%	0.4%	**
Intermountain Medical Center	53	0.0%	1.4%	**
LDS Hospital	86	1.2%	0.8%	**
McKay-Dee Hospital	62	1.6%	1.6%	**
Ogden Regional	31	0.0%	0.8%	**
St. Mark's Hospital	58	0.0%	0.3%	**
University Health Care	134	0.7%	0.8%	**
Utah Valley Regional	83	1.2%	0.8%	**

All eight Utah hospitals in this table had about the same rate of deaths as expected ()
compared to similar patients in the U.S. Half (4 of 8) had no deaths in three years.**

[See additional hospitals that treated at least one patient but less than 30 patients.](#)

[View Data Limitations.](#)

Rating: based on the Exact 95% Confidence Interval test of statistical significance.

*** Lower % deaths than expected, ** Same % deaths as expected, * Higher % deaths than expected

Actual deaths: percentage of patients who received care for this condition or procedure and died in this hospital.

Expected deaths: percentage of patients who were expected to die at this hospital if it performed the same as other U.S. hospitals that treated similar patients (age, gender, how ill the patients were).